






Má-Eö

Distance from the shore (m)	Temperature (°C)
0 m	18.0
250 m	17.5
500 m	17.0
750 m	16.5

Figure 1. Schematic diagram of the experimental setup. The subject is seated in a chair, viewing the screen. The screen displays the target (a red dot) and the starting position (a black dot). The subject's hand is positioned at the starting position. The screen is divided into two regions: a central region (white) and a peripheral region (black). The subject's hand is positioned at the starting position in the central region. The target is located in the peripheral region. The subject's hand is moved towards the target. The screen displays the hand's position and the target's position. The screen is divided into two regions: a central region (white) and a peripheral region (black). The subject's hand is positioned at the starting position in the central region. The target is located in the peripheral region. The subject's hand is moved towards the target. The screen displays the hand's position and the target's position.



Figure 1. Schematic diagram of the experimental setup. The subject is seated in a chair, viewing the screen. The screen displays the target (a red dot) and the starting position (a black dot). The subject is instructed to move the hand from the starting position to the target. The distance between the starting position and the target is 10 cm. The subject is instructed to move the hand at a constant velocity. The screen is 100 cm high and 100 cm wide. The starting position is 50 cm from the screen. The target is 10 cm from the screen. The subject is instructed to move the hand from the starting position to the target. The distance between the starting position and the target is 10 cm. The subject is instructed to move the hand at a constant velocity. The screen is 100 cm high and 100 cm wide. The starting position is 50 cm from the screen. The target is 10 cm from the screen.